



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The Determination of the Conjugate Points for Discontinuous Solutions in the Calculus of Variations.

BY OSKAR BOLZA.

In §§ 8 and 9 of his Inaugural-Dissertation, “*Ueber die discontinuierlichen Lösungen in der Variationsrechnung*” (Göttingen, 1904), Caratheodory develops the general theory of the conjugate points for discontinuous solutions. The object of the present note is to derive Caratheodory’s results concerning conjugate points by a more direct method, to supplement them in certain points, and to give in particular, in explicit form, the equation which connects the parameters of a pair of conjugate points.

§ 1. *Sets of “Broken Extremals”.*

In order that a curve $P_1 P_0 P_2$ with a “corner” at P_0 , but otherwise of class* C' , may minimize† the integral

$$J = \int_{t_1}^{t_2} F(x, y, x', y') dt,$$

it is in the first place necessary that the two “continuous” branches $P_1 P_0$ and $P_0 P_2$ should separately satisfy the four necessary conditions for a minimum with fixed endpoints. In particular, each one of the two arcs $P_1 P_0$ and $P_0 P_2$ must be an *extremal*.

Further, at the point $P_0 (x_0, y_0)$ Weierstrass-Erdmann’s *corner-condition*‡ must be satisfied:

$$\begin{aligned} F_{x'}(x_0, y_0, \cos \mathfrak{D}_0, \sin \mathfrak{D}_0) &= F_{x'}(x_0, y_0, \cos \bar{\mathfrak{D}}_0, \sin \bar{\mathfrak{D}}_0), \\ F_{y'}(x_0, y_0, \cos \mathfrak{D}_0, \sin \mathfrak{D}_0) &= F_{y'}(x_0, y_0, \cos \bar{\mathfrak{D}}_0, \sin \bar{\mathfrak{D}}_0), \end{aligned} \tag{1}$$

* Compare for the terminology my *Lectures on the Calculus of Variations*, § 2, c) and § 24, a).

† In the sense defined in § 24, c) of my *Lectures* and under the assumptions concerning the function $F(x, y, x', y')$ stated in § 24, b).

‡ Compare *Lectures*, § 25, c).

where \mathfrak{S}_0 denotes the amplitude of the positive tangent to the arc $P_1 P_0$ at P_0 , $\bar{\mathfrak{S}}_0$ the amplitude of the positive tangent to the arc $P_0 P_2$ at P_0 .

We shall call a curve $P_1 P_0 P_2$ consisting of two arcs of extremals $P_1 P_0$ and $P_0 P_2$ a “*broken extremal*”, if at P_0 this corner-condition (1) is satisfied.




FIG. 1.

We assume for the following discussion that the curve $P_1 P_0 P_2$ lies in the interior of the domain of continuity R of the function F (compare *Lectures*, § 24, b), and that Legendre's condition is satisfied in the stronger form*

$$F_1 > 0 \quad (2)$$

along each of the two branches $P_1 P_0$ and $P_0 P_2$.

Let now

$$x = \phi(t, a), \quad y = \psi(t, a) \quad (3)$$

be any one-parameter set of extremals which contains the arc $P_1 P_0$ for $a = a_0$, so that the arc $P_1 P_0$ is representable by the equations

$$x = \phi(t, a_0), \quad y = \psi(t, a_0), \quad t_1 \leq t \leq t_0. \quad (4)$$

The functions

$$\phi, \phi_t, \phi_{tt}; \quad \psi, \psi_t, \psi_{tt}$$

* Compare *Lectures*, § 27, b).

are supposed* to be of class C' as functions of t and a in the domain

$$t_1 - h_1 \leq t \leq t_0 + h_0, \quad |a - a_0| \leq d,$$

h_0, h_1, d being sufficiently small positive quantities.

The extremal of the set (3) corresponding to a particular value a will be denoted by \mathfrak{E}_a ; further we write \mathfrak{E}_0 for \mathfrak{E}_{a_0} .

We propose to determine a point $P(t)$ on a given extremal \mathfrak{E}_a of the set (3), and at the same time a direction $\bar{\mathfrak{S}}$ passing through P , such that the direction $\bar{\mathfrak{S}}$ together with the direction \mathfrak{S} of the positive tangent to the extremal \mathfrak{E}_a at P shall satisfy Weierstrass-Erdmann's corner-condition for the point P .

We have, then, for the determination of the two unknown quantities t and $\bar{\mathfrak{S}}$, the two equations†:

$$\begin{aligned} F_{x'}[\phi(t, a), \psi(t, a), \phi_t(t, a), \psi_t(t, a)] - F_{x'}[\phi(t, a), \psi(t, a), \cos \bar{\mathfrak{S}}, \sin \bar{\mathfrak{S}}] &= 0, \\ F_{y'}[\phi(t, a), \psi(t, a), \phi_t(t, a), \psi_t(t, a)] - F_{y'}[\phi(t, a), \psi(t, a), \cos \bar{\mathfrak{S}}, \sin \bar{\mathfrak{S}}] &= 0. \end{aligned} \quad (5)$$

These equations are satisfied for $t = t_0, a = a_0, \bar{\mathfrak{S}} = \bar{\mathfrak{S}}_0$, since according to our assumptions the broken extremal $P_1 P_0 P_2$ satisfies the corner-condition (1). Further, their left-hand members, which we denote by $\Phi(t, a, \bar{\mathfrak{S}})$ and $\Psi(t, a, \bar{\mathfrak{S}})$ respectively, are of class C' in the vicinity of the point $t_0, a_0, \bar{\mathfrak{S}}_0$. Hence we can apply the theorem on implicit functions,‡ provided that the Jacobian

$$J_{t\bar{\mathfrak{S}}} = \frac{\partial(\Phi, \Psi)}{\partial(t, \bar{\mathfrak{S}})}$$

is different from zero at the point $t_0, a_0, \bar{\mathfrak{S}}_0$. If we write for brevity

$$\cos \mathfrak{S} = p, \sin \mathfrak{S} = q; \cos \bar{\mathfrak{S}} = \bar{p}, \sin \bar{\mathfrak{S}} = \bar{q},$$

and remember that along the extremal $P_1 P_0$

$$\frac{\partial}{\partial t} F_{x'} = F_x, \quad \frac{\partial F_{y'}}{\partial t} = F_y,$$

we obtain:

$$\begin{aligned} \Phi_t &= F_x - \bar{F}_{x'x} x' - \bar{F}_{x'y} y', & \Psi_t &= F_y - \bar{F}_{y'x} x' - \bar{F}_{y'y} y', \\ \Phi_{\bar{\mathfrak{S}}} &= \bar{F}_1 \bar{q}, & \Psi_{\bar{\mathfrak{S}}} &= -\bar{F}_1 \bar{p}, \end{aligned} \quad (6)$$

* The existence of an infinitude of sets of extremals satisfying these conditions is a consequence of our assumptions according to certain existence theorems on differential equations; compare Kneser, *Lehrbuch der Variationsrechnung*, § 27, and Bolza, *Trans. Amer. Math. Soc.*, Vol. VII (1906), p. 464.

† Since $F_{x'}, F_{y'}$ are positively homogeneous of dimension zero in x', y' , we may replace in these functions $\cos \mathfrak{S}, \sin \mathfrak{S}$ by $\phi_t(t, a), \psi_t(t, a)$.

‡ Compare, for instance, Osgood, *Lehrbuch der Functionentheorie*, Vol. I, p. 52.

where the arguments of F_x, F_y are: $\phi(t, a), \psi(t, a), x' = \phi_t(t, a), y' = \psi_t(t, a)$; those of $\bar{F}_1, \bar{F}_{x'x}$, etc.: $\phi(t, a), \psi(t, a), \bar{p}, \bar{q}$.

Making use of the homogeneity properties* of the function F and its partial derivatives, we obtain for the above Jacobian:

$$J_{t\bar{s}} = \sqrt{x'^2 + y'^2} \bar{F}_1 \{ p \bar{F}_x + q \bar{F}_y - (\bar{p} F_x + \bar{q} F_y) \}, \quad (7)$$

where now the two last arguments in F_x, F_y are p, q .

The first two factors of $J_{t\bar{s}}$ are different from zero for $t = t_0, a = a_0, \bar{S} = \bar{S}_0$. Hence if we put, with Caratheodory,

$$\begin{aligned} \Omega(x_0, y_0) = & p_0 F_x(x_0, y_0, \bar{p}_0, \bar{q}_0) + q_0 F_y(x_0, y_0, \bar{p}_0, \bar{q}_0) \\ & - \bar{p}_0 F_x(x_0, y_0, p_0, q_0) - \bar{q}_0 F_y(x_0, y_0, p_0, q_0), \end{aligned} \quad (8)$$

where

$$p_0 = \cos \mathfrak{S}_0, \quad q_0 = \sin \mathfrak{S}_0, \quad \bar{p}_0 = \cos \bar{\mathfrak{S}}_0, \quad \bar{q}_0 = \sin \bar{\mathfrak{S}}_0,$$

we have the result:

If the condition

$$\Omega(x_0, y_0) \neq 0 \quad (9)$$

is satisfied, there exists one and but one system of functions

$$t = t(a), \quad \bar{S} = \bar{S}(a), \quad (10)$$

of class C' in the vicinity of $a = a_0$, which satisfies the two equations (5) and the initial conditions

$$t(a_0) = t_0, \quad \bar{S}(a_0) = \bar{S}_0. \quad (11)$$

The functions (10) represent, at least for the vicinity of the point P_0 , the solution of the problem proposed above.

From our assumption (2), applied to the point P_0 and the branch $P_0 P_2$, it follows that

$$F_1(\phi[t(a), a], \psi[t(a), a], \cos \mathfrak{S}(a), \sin \bar{\mathfrak{S}}(a)) \neq 0$$

for all sufficiently small values of $|a - a_0|$. Hence† it is possible to construct one and but one extremal

$$\bar{\mathfrak{C}}_a: \quad x = \bar{\phi}(t, a), \quad y = \bar{\psi}(t, a) \quad (12)$$

through the point P in the direction $\mathfrak{S}(a)$. The parameter t can be so selected that also on $\bar{\mathfrak{C}}_a$ the value $t = t(a)$ furnishes the point P , so that

$$\bar{\phi}[t(a), a] = \phi[t(a), a], \quad \bar{\psi}[t(a), a] = \psi[t(a), a]. \quad (13)$$

* Compare *Lectures*, §24, b) equations (8) and (10).

† According to Cauchy's existence theorem on differential equations; compare *Lectures*, §25, b).

We thus obtain a broken extremal $\mathfrak{E}_a + \bar{\mathfrak{E}}_a$ with a corner at P , on which the parameter t varies continuously. If we let a vary, we obtain a set of broken extremals. We shall call the set (12) the set of extremals complementary to the set (3). On account of (11) it contains, for $a = a_0$, the extremal $\bar{\mathfrak{E}}_0$ of which the arc $P_0 P_2$ forms a part.

From the properties of the integrals of a system of differential equations as functions of their initial values,* it follows that the functions $\bar{\phi}(t, a)$, $\bar{\psi}(t, a)$ have the same continuity properties as the functions $\phi(t, a)$, $\psi(t, a)$, in a domain

$$t_0 - \bar{h}_0 \leq t \leq t_2 + \bar{h}_2, \quad |a - a_0| \leq \bar{d}.$$

§ 2. The Corner-Curve.†

If we let a vary, the corner P describes a curve $\tilde{\mathfrak{C}}$, which we call the "corner-curve". If we define the functions $\tilde{x}(a)$, $\tilde{y}(a)$ by the equations

$$\tilde{x}(a) = \phi[t(a), a], \quad \tilde{y}(a) = \psi[t(a), a], \quad (14)$$

or, what amounts to the same thing according to (13),

$$\tilde{x}(a) = \bar{\phi}[t(a), a], \quad \tilde{y}(a) = \bar{\psi}[t(a), a], \quad (14a)$$

the corner-curve is given in parameter-representation by the equations

$$\tilde{\mathfrak{C}}: \quad x = \tilde{x}(a), \quad y = \tilde{y}(a),$$

and any particular value of a furnishes that point of $\tilde{\mathfrak{C}}$ which is the corner for the corresponding broken extremal $\mathfrak{E}_a + \bar{\mathfrak{E}}_a$.

We propose first to compute the slope $\tan \tilde{\mathfrak{S}}$ of the tangent to the corner-curve $\tilde{\mathfrak{C}}$ at the point P .

From the definition of the functions \tilde{x} , \tilde{y} , we obtain for their derivatives with respect to a :

$$\tilde{x}' = \phi_t t'(a) + \phi_a, \quad \tilde{y}' = \psi_t t'(a) + \psi_a;$$

and from (5) we obtain, according to the rules for the differentiation of implicit functions,

$$t'(a) = - \frac{J_{a\bar{\partial}}}{J_{t\bar{\partial}}},$$

where

$$J_{a\bar{\partial}} = \frac{\partial(\Phi, \Psi)}{\partial(a, \mathfrak{S})}.$$

* Compare Kneser, *Lehrbuch der Variationsrechnung*, §27, and Bliss, The Solution of Differential Equations of the First Order as Functions of their Initial Values, *Annals of Mathematics*, Ser. 2, Vol. VI, p. 49.

† Caratheodory's "Knickpunkt-Curve".

But

$$\begin{aligned}\Phi_a &= F_{x'x} \phi_a + F_{x'y} \psi_a + F_{x'x'} \phi_{ta} + F_{x'y'} \psi_{ta} - \bar{F}_{x'x} \phi_a - \bar{F}_{x'y} \psi_a, \\ \Psi_a &= F_{y'x} \phi_a + F_{y'y} \psi_a + F_{y'x'} \phi_{ta} + F_{y'y'} \psi_{ta} - \bar{F}_{y'x} \phi_a - \bar{F}_{y'y} \psi_a;\end{aligned}$$

the functions $\bar{F}_{x'x}$, $\bar{F}_{x'y}$, $\bar{F}_{y'x}$, $\bar{F}_{y'y}$ are positively homogeneous of dimension zero with respect to their last two arguments \bar{p} , \bar{q} ; hence we may replace \bar{p} and \bar{q} by $\phi_t(t, a)$ and $\psi_t(t, a)$ respectively. This being done, we express all the partial derivatives of F in terms of Weierstrass' functions*: F_1, L, M, N . The result is

$$\begin{aligned}\Phi_a &= -A \phi_a - B \psi_a - y' \Delta_t F_1 - \bar{y}' \bar{F}_1 (\phi_a \bar{y}'' - \psi_a \bar{x}''), \\ \Psi_a &= -B \phi_a - C \psi_a + x' \Delta_t F_1 + \bar{x}' \bar{F}_1 (\phi_a \bar{y}'' - \psi_a \bar{x}''),\end{aligned}\tag{15}$$

where

$$\begin{aligned}x' &= \phi_t(t, a), \quad y' = \psi_t(t, a); \quad \bar{x}' = \bar{\phi}_t(t, a), \quad \bar{y}' = \bar{\psi}_t(t, a); \\ \bar{x}'' &= \bar{\phi}_{tt}(t, a), \quad \bar{y}'' = \bar{\psi}_{tt}(t, a); \quad \Delta(t, a) = \phi_t \psi_a - \psi_t \phi_a, \\ A &= \bar{L} - L, \quad B = \bar{M} - M, \quad C = \bar{N} - N;\end{aligned}$$

the quantities L, M, N refer to the point P and the extremal \mathfrak{E}_a , the quantities $\bar{L}, \bar{M}, \bar{N}$ to the point P and the extremal $\bar{\mathfrak{E}}_a$. Finally, the last two arguments of F_1 and \bar{F}_1 are x', y' and \bar{x}', \bar{y}' respectively.

From (15) and (6) we obtain

$$J_{a\bar{a}} = (\bar{x}'^2 + \bar{y}'^2) \bar{F}_1 \{ \phi_a (A \bar{x}' + B \bar{y}') + \psi_a (B \bar{x}' + C \bar{y}') - \Delta_t F_1 (x' \bar{y}' - y' \bar{x}') \}.\tag{16}$$

At the same time the expression (7) for $J_{t\bar{a}}$ may be thrown into another form, if we remember the homogeneity properties of F_1, F_x, F_y and make use of the relations†

$$Lx' + My' = F_x, \quad Mx' + Ny' = F_y;$$

we thus obtain

$$J_{t\bar{a}} = (\bar{x}'^2 + \bar{y}'^2) \bar{F}_1 [A x' \bar{x}' + B (x' \bar{y}' + y' \bar{x}') + C y' \bar{y}'].\tag{17}$$

The comparison between the two expressions for $J_{t\bar{a}}$ leads to a second form for the quantity $\Omega(x, y)$; viz.,

$$\Omega(x, y) = A p \bar{p} + B (p \bar{q} + q \bar{p}) + C q \bar{q}.\tag{18}$$

We thus finally obtain

$$\begin{aligned}\bar{x}' &= -\frac{\Delta(B \bar{x}' + C \bar{y}') + x' \Delta_t F_1 (x' \bar{y}' - y' \bar{x}')}{A x' \bar{x}' + B (x' \bar{y}' + y' \bar{x}') + C y' \bar{y}'}, \\ \bar{y}' &= \frac{\Delta(A \bar{x}' + B \bar{y}') + y' \Delta_t F_1 (x' \bar{y}' - y' \bar{x}')}{A x' \bar{x}' + B (x' \bar{y}' + y' \bar{x}') + C y' \bar{y}'}\end{aligned}\tag{19}$$

* Compare *Lectures*, Chap. IV, equations (11 a) and (35).

† Compare *Lectures*, p. 132.

Hence follows, for the slope $\tan \tilde{\mathfrak{S}}$ of the tangent to the corner-curve $\tilde{\mathfrak{C}}$ at the point P , the expression

$$\tan \tilde{\mathfrak{S}} = \frac{\Delta (A \bar{x}' + B \bar{y}') + y' \Delta_t F_1 (x' \bar{y}' - y' \bar{x}')}{-\Delta (B \bar{x}' + C \bar{y}') + x' \Delta_t F_1 (x' \bar{y}' - y' \bar{x}')} . \quad (20)$$

§ 3. *Interrelation Between the Slope of the Corner-Curve at P_0 and the Focal-Points of the Set of Broken Extremals.*

We now consider in particular the question how the slope $\tan \tilde{\mathfrak{S}}_0$ of the tangent to the corner-curve at P_0 depends upon the choice of the set of extremals (3). For this purpose we have to put $\alpha = \alpha_0$ in (20), and consequently, according to (11), the argument $t = t(\alpha)$, in $x', y'; \bar{x}', \bar{y}', \Delta(t, \alpha)$, etc., equal to t_0 . In the resulting expression for $\tan \tilde{\mathfrak{S}}_0$, the Jacobian $\Delta(t_0, \alpha_0)$ and its derivative $\Delta_t(t_0, \alpha_0)$ are the only quantities which depend upon the choice of the set of extremals (3).

The function $\Delta(t, \alpha_0)$, in its turn, is determined to a constant factor by the condition that it satisfies Jacobi's differential equation* for the extremal \mathfrak{C}_0 , viz.,

$$F_2 u - \frac{d}{dt} \left(F_1 \frac{du}{dt} \right) = 0, \quad (21)$$

and by one of its zeros. Let $t = \tau$ be the zero of $\Delta(t, \alpha_0)$ next smaller than t_0 , so that the corresponding point of \mathfrak{C}_0 , which we denote by Q , is the focal point† of the set (3) on \mathfrak{C}_0 . Then

$$\Delta(t, \alpha_0) = \text{Const. } \Theta(t, \tau),$$

where $\Theta(t, \tau)$ is the function which determines in Weierstrass'‡ theory the conjugate point to Q . We may therefore write

$$\tan \tilde{\mathfrak{S}}_0 = \frac{\alpha \Theta(t_0, \tau) + \beta \Theta_t(t_0, \tau)}{\gamma \Theta(t_0, \tau) + \delta \Theta_t(t_0, \tau)}, \quad (22)$$

where

$$\begin{aligned} \alpha &= A_0 \bar{p}_0 + B_0 \bar{q}_0, & \beta &= q_0 F_1(t_0) \sin(\mathfrak{S}_0 - \mathfrak{S}_0)(x_0'^2 + y_0'^2), \\ \gamma &= -(B_0 \bar{p}_0 + C_0 \bar{q}_0), & \delta &= p_0 F_1(t_0) \sin(\tilde{\mathfrak{S}}_0 - \mathfrak{S}_0)(x_0'^2 + y_0'^2), \end{aligned} \quad (23)$$

the subscript 0 indicating that the quantities to which it is affixed are to be computed for the point P_0 .

* Compare *Lectures*, pp. 40 and 200.

† Compare Kneser, *Lehrbuch der Variationsrechnung*, § 24, and my *Lectures*, § 38.

‡ Compare *Lectures*, p. 135.

The coefficients $\alpha, \beta, \gamma, \delta$ are therefore independent of τ . Hence the slope of the corner-curve $\tilde{\mathfrak{C}}$ at P_0 is the same for all sets of extremals (3) which have the same focal point Q , the set of extremals through the point Q being included among the latter.

We examine next how the slope $\tan \tilde{\mathfrak{S}}_0$ varies when the focal point Q describes the extremal \mathfrak{E}_0 . For this purpose, we compute the derivative of $\tan \tilde{\mathfrak{S}}_0$ with respect to τ :

$$\frac{d \tan \tilde{\mathfrak{S}}_0}{d\tau} = - \frac{(\alpha \delta - \beta \gamma) \{ \Theta(t_0, \tau) \Theta_{t\tau}(t_0, \tau) - \Theta_t(t_0, \tau) \Theta_\tau(t_0, \tau) \}}{\{ \gamma \Theta(t_0, \tau) + \delta \Theta_t(t_0, \tau) \}^2}.$$

But from the definition of $\Theta(t, \tau)$ it follows that

$$\begin{aligned} \Theta(t_0, \tau) \Theta_{t\tau}(t_0, \tau) - \Theta_t(t_0, \tau) \Theta_\tau(t_0, \tau) \\ = [\mathfrak{S}_1(t_0) \mathfrak{S}'_2(t_0) - \mathfrak{S}_2(t_0) \mathfrak{S}'_1(t_0)] [\mathfrak{S}_1(\tau) \mathfrak{S}'_2(\tau) - \mathfrak{S}_2(\tau) \mathfrak{S}'_1(\tau)], \end{aligned}$$

where $\mathfrak{S}_1(t), \mathfrak{S}_2(t)$ are two linearly independent solutions of Jacobi's differential equation (21). Hence from the theory of linear differential equations it follows* that

$$\mathfrak{S}_1(t) \mathfrak{S}'_2(t) - \mathfrak{S}_2(t) \mathfrak{S}'_1(t) = \frac{k}{F_1(t)},$$

where k is a constant different from zero.

On the other hand we get, on substituting the values of $\alpha, \beta, \gamma, \delta$,

$$\alpha \delta - \beta \gamma = F_1(t_0) \sin(\mathfrak{S}_0 - \mathfrak{S}_0) \Omega(x_0, y_0) (x_0'^2 + y_0'^2).$$

Hence it follows that

$$\frac{d}{d\tau} \tan \tilde{\mathfrak{S}}_0 = \frac{-k^2 (x_0'^2 + y_0'^2) \sin(\mathfrak{S}_0 - \mathfrak{S}_0) \Omega(x_0, y_0)}{F_1(\tau) \{ \gamma \Theta(t_0, \tau) + \delta \Theta_t(t_0, \tau) \}^2}. \quad (24)$$

We suppose for the further discussion that

$$\mathfrak{S}_0 - \mathfrak{S}_0 \not\equiv 0 \pmod{\pi}, \quad (25)$$

and that the inequality (2) holds not only for the arc $P_1 P_0$ of the extremal \mathfrak{E}_0 but also for its continuation beyond P_1 , at least as far as the point $P'_0 (t = t'_0)$ whose conjugate the point P_0 is.

And now we let τ increase from t'_0 to t_0 ; i. e., we let the point Q describe the extremal \mathfrak{E}_0 from P'_0 to P_0 . The derivative of $\tan \tilde{\mathfrak{S}}_0$ will then always have a

* Compare, for instance, *Lectures*, p. 58, footnote².

constant sign, since $\Omega(x_0, y_0)$, which is independent of τ , is supposed to be different from zero. For $\tau = t'_0$ and $\tau = t_0$, but for no other value between them, the function $\Theta(t_0, \tau)$ vanishes and $\tan \tilde{S}_0$ takes the value

$$\tan \tilde{S}_0 = \frac{\beta}{\delta} = \frac{q_0}{p_0} = \tan S_0.$$

Hence we have the result:

While the point Q describes the extremal \mathfrak{G}_0 from P'_0 to P_0 , the line \tilde{S}_0 revolves about the point P_0 from the initial position S_0 constantly in the same sense through an angle of 180° . The rotation takes place:*

In positive sense, when $\Omega(x_0, y_0) \sin(\bar{S}_0 - S_0) < 0$;

In negative sense, when $\Omega(x_0, y_0) \sin(\bar{S}_0 - S_0) > 0$.

It passes therefore once and but once through the position \tilde{S}_0 . We denote the value of τ for which this takes place by e_0 and the corresponding point† of \mathfrak{G}_0 by E_0 . For the discussion of sufficient conditions, it is important to distinguish whether the line \tilde{S}_0 lies in the angle‡ between the two branches $P_1 P_0$ and $P_0 P_2$ or outside of it. Four cases must be distinguished according to the signs of $\Omega(x_0, y_0)$ and $\sin(\bar{S}_0 - S_0)$. The result is:

While the point Q moves from P'_0 to E_0 , the line \tilde{S}_0 revolves from the position S_0 into the position \tilde{S}_0 , inside of the angle between $P_1 P_0$ and $P_0 P_2$ when $\Omega(x_0, y_0) > 0$, outside of it when $\Omega(x_0, y_0) < 0$. As the point Q moves on from E_0 to P_0 , the line \tilde{S}_0 continues its rotation from the position \tilde{S}_0 into the position S_0 , outside of the angle in question when $\Omega(x_0, y_0) > 0$, inside of it when $\Omega(x_0, y_0) < 0$.

Conversely: To every line \tilde{S}_0 through the point P_0 which is tangent to neither of the two arcs $P_1 P_0$, $P_0 P_2$ at P_0 , there belongs one and but one point Q , between P'_0 and P_0 , such that the corner-curve for every set of extremals (3) for which Q is the focal point, touches the line \tilde{S}_0 at P_0 .

The value of τ belonging to a given line \tilde{S}_0 is obtained by solving equation (22) with respect to τ . The equation may be thrown into the form

$$\begin{aligned} & [A_0 \bar{p}_0 \tilde{p}_0 + B_0 (\bar{p}_0 \tilde{q}_0 + \bar{q}_0 \tilde{p}_0) + C_0 \bar{q}_0 \tilde{q}_0] \Theta(t_0, \tau) \\ & - (x_0'^2 + y_0'^2) F_1(t_0) \sin(\bar{S}_0 - S_0) \sin(\tilde{S}_0 - S_0) \Theta_t(t_0, \tau) = 0, \end{aligned} \quad (26)$$

where

$$\tilde{p}_0 = \cos \tilde{S}_0, \quad \tilde{q}_0 = \sin \tilde{S}_0.$$

* I. e., the line through P_0 of slope $\tan \tilde{S}_0$.

† Caratheodory denotes this point by E_1 ; see *Dissertation*, p. 31.

‡ I. e., that one of the two angles formed by the half-rays $\bar{\vartheta}_0$ and $\vartheta_0 + \pi$ which is less than π .

Case I: $\sin(\bar{\vartheta}_0 - \vartheta_0) > 0$, $\Omega(x_0, y_0) > 0$.




FIG. 2.

Case II: $\sin(\bar{\vartheta}_0 - \vartheta_0) > 0$, $\Omega(x_0, y_0) < 0$.




FIG. 3.

Case III: $\sin(\bar{\vartheta}_0 - \vartheta_0) < 0$, $\Omega(x_0, y_0) > 0$.




FIG. 4.

Case IV: $\sin(\bar{\vartheta}_0 - \vartheta_0) < 0$, $\Omega(x_0, y_0) < 0$.




FIG. 5.

In particular, the equation for the determination of the parameter e_0 of the point E_0 is obtained by putting, in (26), $\bar{S}_0 = S_0$.

§ 4. *The Conjugate Points of Discontinuous Solutions.*

Let, for a moment, the equations (12) represent any set of extremals containing, for $a = a_0$, the extremal $\bar{\mathfrak{E}}_0$. We may then propose for the set (12) the same problem which we have solved in § 1 for the set (3). The only difference will be that in the equations (5) the symbols ϕ, ψ, \bar{S} must be interchanged with $\bar{\phi}, \bar{\psi}, S$, and the same interchange must be applied in the results; in this process the quantities A, B, C are changed into $-A, -B, -C$. Accordingly, if $\bar{Q}(t = \bar{\tau})$ be the focal point of the set (12) on $\bar{\mathfrak{E}}_0$, the slope of the corner-curve belonging to the set (12) at P_0 is

$$\tan \bar{S}_0 = \frac{\bar{\alpha} \bar{\Theta}(t_0, \bar{\tau}) + \bar{\beta} \bar{\Theta}_t(t_0, \bar{\tau})}{\bar{\gamma} \bar{\Theta}(t_0, \bar{\tau}) + \bar{\delta} \bar{\Theta}_t(t_0, \bar{\tau})}, \quad (27)$$

where the quantities $\bar{\alpha}, \bar{\beta}, \bar{\gamma}, \bar{\delta}$ are derived from $\alpha, \beta, \gamma, \delta$ by the above interchange, and $\bar{\Theta}$ has the same meaning for $\bar{\mathfrak{E}}_0$ as Θ for \mathfrak{E}_0 .

Conversely, we obtain the value of $\bar{\tau}$ corresponding to a given line \bar{S} by solving equation (27). We denote the value of $\bar{\tau}$ corresponding to the particular line S_0 by \bar{e}_0 and the corresponding point of $\bar{\mathfrak{E}}_0$ by \bar{E}_0 ; this point lies between the point P_0 and its conjugate $\bar{P}'_0(t = \bar{t}'_0)$ on $\bar{\mathfrak{E}}_0$.

Let now the equation (12) denote again, as in § 1, the particular set of extremals complementary to the set (3). The two sets (3) and (12) will then have the corner-curve in common; hence we have, in this case,

$$\bar{S}_0 = S_0.$$

We obtain, therefore, the focal point of the set (12) complementary to the set (3) by equating the right-hand members of the two equations (22) and (27) and solving the equation thus obtained with respect to τ . After some reductions the following result is obtained:

If $t = \tau$ be the parameter of the focal point Q of the set of extremals (3) on \mathfrak{E}_0 , and $t = \bar{\tau}$ the parameter of the focal point Q of the set (12), complementary to (3), on $\bar{\mathfrak{E}}_0$, then the following relation holds between τ and $\bar{\tau}$:

$$\left. \begin{aligned}
 & (A_0 C_0 - B_0^2) \Theta(t_0, \tau) \bar{\Theta}(t_0, \bar{\tau}) \\
 & - (x_0'^2 + y_0'^2) F_1(t_0) (A_0 p_0^2 + 2 B_0 p_0 q_0 + C_0 q_0^2) \frac{\partial \Theta(t_0, \tau)}{\partial t_0} \bar{\Theta}(t_0, \tau) \\
 & + (\bar{x}_0'^2 + \bar{y}_0'^2) \bar{F}_1(t_0) (A_0 \bar{p}_0^2 + 2 B_0 \bar{p}_0 \bar{q}_0 + C_0 \bar{q}_0^2) \Theta(t_0, \tau) \frac{\partial \bar{\Theta}(t_0, \bar{\tau})}{\partial t_0} \\
 & - (x_0'^2 + y_0'^2) (\bar{x}_0'^2 + \bar{y}_0'^2) F_1(t_0) \bar{F}_1(t_0) \sin^2(\bar{\mathfrak{S}}_0 - \mathfrak{S}_0) \frac{\partial \Theta(t_0, \tau)}{\partial t_0} \frac{\partial \bar{\Theta}(t_0, \bar{\tau})}{\partial t_0} = 0.
 \end{aligned} \right\} \quad (28)$$

The two points Q and \bar{Q} are called, according to Caratheodory,* a pair of conjugate points of the broken extremal $\mathfrak{E}_0 + \bar{\mathfrak{E}}_0$. According to a previous remark, the point \bar{Q} conjugate to Q on $\mathfrak{E}_0 + \bar{\mathfrak{E}}_0$ may also be defined as the focal point on $\bar{\mathfrak{E}}_0$ of the set of extremals which is complementary to the set of extremals through the point Q .

In Figs. 2 to 5 the interrelation between the points Q and \bar{Q} and the line $\tilde{\mathfrak{S}}_0$ is indicated. For instance, in Case I the point \bar{Q} moves on $\bar{\mathfrak{E}}_0$ from \bar{E}_0 to \bar{P}'_0 while the point Q moves on \mathfrak{E}_0 from P'_0 to E_0 ; at the same time the line $\tilde{\mathfrak{S}}_0$ revolves about P_0 from the position \mathfrak{S}_0 , in the sense of the arrow, into the position $\bar{\mathfrak{S}}_0$.

The conjugate points thus defined play for the discontinuous solutions a rôle similar to that of the ordinary conjugate points for continuous solutions, at least in the case when the line $\tilde{\mathfrak{S}}_0$ lies inside the angle of the two branches $P_1 P_0, P_0 P_2$. We refer in this respect to Caratheodory's dissertation, § 9.

THE UNIVERSITY OF CHICAGO, January 29, 1907.

* Caratheodory restricts, however, the definition to the case when the line $\tilde{\mathfrak{S}}_0$ lies inside of the angle of the two branches $P_1 P_0, P_0 P_2$.